

# Presentation and outcomes of neonatal sepsis at Beni-Suef university hospital

*By Reda Reyad*

## Presentation and outcomes of neonatal sepsis at Beni-Suef university hospital

Mona Nasser<sup>1</sup>, Reda Reyad<sup>2</sup>, Sameh Fahmy<sup>2</sup>, Amna Mabrouk<sup>2</sup>

5

<sup>1</sup>Clinical and Chemical Pathology Department, Faculty of Medicine, Beni-Suef University, Egypt

<sup>2</sup>Pediatrics Department, Faculty of medicine, Beni-Suef University, Egypt

[redaokasha88@gmail.com](mailto:redaokasha88@gmail.com)

Reda Reyad

11

### ABSTRACT

**Background.** Sepsis is the third leading cause of neonatal death, especially in developing countries, and it is characterized by severe infections caused by many pathogens and subsequent organ failure.

**Aim of the work.** Our study set out to identify the signs, symptoms, and outcomes of neonatal sepsis in both the clinic and the laboratory of university hospital of Beni-Suef.

**Methodology.** The pediatric department at Beni-Suef University hospital conducted a prospective observational study. One hundred neonates with sepsis were part of the study. In addition to standard laboratory testing (CBC, CRP, and blood culture), all patients had a thorough clinical examination, history recording, and monitoring for complications and outcomes (i.e., improvement or death).

**Results.** One hundred people, with an average gestational age of  $35.4 \pm 2.5$  weeks, were part of the study that was diagnosed with neonatal sepsis. At birth, they weighed an average of  $2.6 \pm 0.7$  kg. The emergency room sent 10 patients, the obstetrics and gynecology department sent 35, and other hospitals sent 55. We determined that 76 of the total cases were late-onset neonatal sepsis and 24 were early-onset. Newborn respiratory distress accounted for 81 cases, making it the leading cause of hospitalization. The most common adverse event was septic shock. Half of the patients showed signs of improvement. Nearly all of the patients (95%) obtained positive results from blood cultures, with the majority of those positive results being for Gram-negative bacteria.

With 46% of the total, Klebsiella was the most common kind of isolated bacteria. There was significant association between mortality and younger gestational age  $P$  value<0.001, female gender  $P$  value 0.015, need to mechanical ventilation  $P$  value<0.001, intravenous infusion of positive inotropic drugs  $p$  value <0.001, need to platelet transfusion  $P$  value<0.001, Complications  $P$  value<0.001, low birth weight  $P$  value0.001 and lower systolic and diastolic blood pressure  $P$  value0.020, 0.009

**Conclusions.** The most common cause for admitting neonates to Beni Suef University Hospital with neonatal sepsis was respiratory distress. Babies born by cesarean section more susceptible for sepsis than vaginal births. The majority of septic neonates were referred to our unit from other hospital while Klebsiella was the most often diagnosed bacterium. Out of 100 neonates who included in our study, 49 neonates were died. Prematurity, low birth weight, female gender, mechanical ventilation, platelet transfusion, intravenous infusion of positive inotropic drugs, complications in neonatal intensive care during staying, and lower blood pressure were considered risk factors for death. AS there were significant associations between these parameters and mortality

**Key words:** neonatal sepsis, mortality, complications, blood culture

## INTRODUCTION

Bacteria, viruses, or fungi may cause neonatal sepsis, a systemic disease. It can cause hemodynamic changes and other symptoms, leading to significant harm or even death. We can classify neonatal sepsis as early-onset, late-onset, or very late-onset based on when symptoms first appear. Babies with sepsis detected beyond the first week of life or between days 4 and 30 are considered to have late-onset neonatal sepsis. When signs and symptoms of newborn sepsis appear during the first three days of a baby's life, rather than after 72 hours, we say that the condition has an early beginning [1]. Over 2.6 million newborns die every year, with 75% of those deaths happening in the first week, according to epidemiological studies [2]. Organisms that enter the body through the mother's vaginal canal or the hospital delivery setting often cause early-onset sepsis [3]. Typical symptom of newborns with early-onset sepsis (EOS) is respiratory distress, which is associated with high rates of morbidity and mortality [4]. Pathogens brought into the hospital by patients or their caregivers are a common cause of Late onset sepsis [5]. While there has been an uptick in the infant survival rate in neonatal intensive care units (NICU) recently, there has also been an increase in the risks of hospital-acquired infections, a leading cause of neonatal mortality. Severe systemic infections affect over 20% of premature and low birth weight infants admitted to the NICU [6]. Averting disastrous results may be possible with early diagnosis and swift treatment.

## PATIENT AND METHODS

### Patients

Prospective observational study was conducted at pediatric department at Beni-Suef university Hospital. The study included 100 patients with neonatal sepsis, 63 male and 37 female, 76 neonate was delivered by cesarean section and 24 was delivered by normal vaginal delivery.

Patients must meet the following diagnostic criteria for infant sepsis in order to participate in the experiment: A baby has sepsis if a blood culture comes back positive and there are other signs of infection, like a rise in C-reactive protein, a drop in white blood cell count (leucopenia or leucocytosis), and a drop in platelet count (thrombocytopenia). In a clinical setting, one may observe symptoms such as temperature changes, low blood pressure, prolonged capillary refill time, irregular heartbeats (tachy or brady), difficulty breathing, restlessness, lack of energy, seizures, poor dietary intake, nausea, vomiting, abdominal distention, yellowing of the skin, abnormal blood vessels, and bleeding.

The following conditions are not eligible for inclusion: severe birth defects, hypoxia, liver disease, kidney failure, cancer, certain drugs (such as linezolid or epinephrine), diabetes mellitus, or metabolic problems that are present at birth.

### Ethical considerations:

Ethical considerations: This study received approval from the study Ethics Committee of the Faculty of Medicine, Beni-Suef University, in September 2020. Informed permission was obtained from the guardians of all study participants before to the investigation.

## METHODS

### All participations are subject to:

**Comprehensive history acquisition:** Maternal medical history: Maternal history of infection during gestation or parturition (kind and duration of antimicrobial treatment), urinary tract infection. Chorioamnionitis, protracted rupture of membranes, difficult delivery, and consanguinity. Medical history of the patient: Gestational age, birth weight, prior NICU admission history, age at admission, reason for NICU hospitalization, and onset timing of sepsis (early or late). Healthcare intervention: Respiratory assistance (nasal, CPAP, mechanical breathing), need for packed red blood cell transfusion, platelet transfusion, or plasma transfusion, necessity for inotropic agents, and length of hospital stay.

**21 Comprehensive clinical evaluation:** Birth weight, gestational age per Dobowitz score, vital signs (heart rate, respiratory rate, temperature, blood pressure), neonatal reflexes such as Moro and suckling, and thorough assessment of all systems including cardiac, respiratory, and neurological (hypotonia, lethargy, seizures, irritability), as well as abdominal examination.

Monitoring of newborn **2** during hospitalization

**Laboratory analyses:** Complete blood count, C-reactive protein, blood culture

**Specimen collection:**

Under aseptic circumstances, blood samples were collected from a peripheral vein in order to conduct CRP, CBC, and blood culture tests. The blood sample size ranged from around 3.5 to 4 ml, with 1 ml set aside for CRP analysis, 1.5 to 2 ml for blood culture **26**, and 1 ml for CBC evaluation. We promptly injected 1.5–2 ml of blood into vials of blood culture medium and submitted them to the clinical microbiology laboratory at our hospital to be cultured and processed further.

#### Methods for Statistics:

SPSS (version 22) and Microsoft Excel (Microsoft Corporation, NY, USA) were used for all statistical analyses. An unpaired t-test was used for inferential research using quantitative variables in two dependent groups with parametric data and the Chi-Square test (X<sup>2</sup>) for comparisons involving categorical data. Two independent samples were compared using the Mann-Whitney test, whereas three or more independent samples were compared using the Kruskal-Wallis test. Odds ratios (OR) with accompanying 95% CIs were used to depict associations. Statistical significance was defined as a p-value less than 0.05.

## RESULTS

Prospective observational study **was** conducted at pediatric department **19** Beni-Suef university Hospital. The study included 100 patients **22** with neonatal sepsis, mean±SD for gestational age, birth weight, was 35.4±2.5 weeks, 2.6±0.7 kg respectively, 63.0% of patient group were males and 76.0% of them were born by cesarean section

**Table (1) Maternal risk factors among the studied patients**

Characteristics	Cases (no=100)	
	No.	%
Urinary tract infection	0.0	0.0
Chorioamnionitis	2	2.0

<b>Premature rupture of membranes</b>	14	14.0
<b>Prenatal hemorrhage</b>	9	9.0
<b>Preeclampsia</b>	7	7.0
<b>Diabetes mellitus</b>	4	4.0
<b>obstructed labor</b>	5	5.0
<b>Hypertension</b>	11	11.0
<b>Polyhydramnios</b>	22	22.0

This table showed no reported cases with urinary tract infection, only 2% with chorioamnionitis, only 14% with premature rupture of membranes

**Table (2) Neonatal history and risk factors among the studied patients:**

<b>Characteristics</b>	<b>Cases (no=100)</b>	
	<b>no.</b>	<b>%</b>
<b>Referral from other hospitals</b>	55	55.0
<b>Emergency department</b>	10	10.0
<b>Gynecology &amp;obstetric department</b>	35	35
<b>Onset of sepsis</b>		
<b>Early</b>	24	24.0
<b>Late</b>	76	76.0
<b>Cause of hospital admission</b>		
<b>Abdominal distension</b>	1	1.0
<b>Convulsions</b>	3	3.0
<b>Encephalocele&amp; respiratory distress</b>	1	1.0
<b>Jaundice</b>	4	4.0
<b>Multiple abscesses &amp; respiratory distress</b>	1	1.0
<b>Poor feeding</b>	1	1.0
<b>respiratory distress only</b>	81	81.0
<b>respiratory distress &amp;convulsions</b>	2	2.0
<b>respiratory distress &amp;gangrene</b>	1	1.0
<b>respiratory distress &amp;Fever</b>	1	1.0
<b>respiratory distress &amp;poor feeding</b>	3	3.0
<b>Vomiting&amp; diarrhea</b>	1	1.0
<b>subarachnoid hemorrhage</b>	1	1.0

<b>Head trauma</b>	1	1.0
<b>Respiratory support</b>		
<b>Nasal oxygen</b>	16	16.0
<b>Continuous positive airway pressure</b>	28	28.0
<b>Mechanical ventilation</b>	54	54.0
<b>Blood elements transfusion</b>		
<b>Packed red blood cells</b>	83	83.0
<b>Plasma</b>	65	65.0
<b>PLT</b>	39	39.0
<b>Need to positive inotropes</b>	57	57.0

This table showed 76% had late onset sepsis, and the most common cause of admission was respiratory distress either alone or with other causes.

**Table (3) General and systemic examination among the studied patients at admission**

Examination		Cases (no=100)	
		no.	%
<b>Birth weight</b>		2.6±0.7	
<b>vital signs</b>	<b>Respiratory rate (mean±SD)</b>	57±18	
	<b>Heart rate(mean±SD)</b>	133±24	
	<b>Temperature(mean±SD)</b>	36.9±0.6	
	<b>systolic blood pressure</b>	73.4±8.5	
	<b>Diastolic blood pressure</b>	43.9±7.5	
<b>Chest</b>	<b>Apnea</b>	3	3.0
	<b>Diminished air entry</b>	30	30.0
	<b>fine crepitation</b>	38	38.0
	<b>fine crepitations &amp; wheezes</b>	12	12.0
	<b>wheezes</b>	3	3.0
	<b>Reparatory distress</b>	88	88.0
<b>Gastro intestinal tract</b>	<b>Poor feeding</b>	73	73.0
	<b>Vomiting</b>	19	19.0
	<b>Distention</b>	46	46.0
	<b>Jaundice</b>	66	66.0
	<b>Bleeding</b>	63	63.0
<b>central nervous system</b>	<b>Irritability</b>	5	5.0
	<b>Lethargy</b>	9	9.0
	<b>Seizures</b>	12	12.0
	<b>Moro reflex</b>		
	<b>Fair</b>	20	20.0
	<b>Good</b>	25	25.0
	<b>Poor</b>	55	55.0
	<b>Suckling reflex</b>		
<b>Fair</b>	19	19.0	

	<b>Good</b>	23	23.0
	<b>Poor</b>	58	58.0

14

This table showed that the mean heart rate, and respiratory rate, systolic blood pressure, diastolic blood pressure was  $133\pm 24$  b/min, and  $57\pm 18$  breath/minute,  $73.4\pm 8.5$ mmHg,  $43.9\pm 7.5$ mmHg respectively and the most common symptom was respiratory distress

**Table (4) outcomes of the studied patients:**

Outcomes	Cases (no=100) N (%)	
<b>Complications of neonatal sepsis</b>		
Acute kidney injury	1	1.0
Central nervous system infection	7	7.0
Cholestasis	1	1.0
Cholestasis and Central nervous system infection	1	1.0
Necrotizing enterocolitis and Pulmonary hemorrhage	1	1.0
Necrotizing enterocolitis and septic shock	1	1.0
Necrotizing enterocolitis and septic shock and acute kidney injury	1	1.0
Pneumothorax	1	1.0
Pneumothorax and Central nervous system infection	1	1.0
Pneumothorax and Necrotizing enterocolitis	1	1.0
Pneumothorax and Septic shock	2	2.0
Pulmonary hemorrhage	6	6.0
Pulmonary hemorrhage and acute kidney injury	2	2.0
Pulmonary hemorrhage and Septic shock	10	10.0
Pulmonary hemorrhage and acute kidney injury	1	1.0



<b>Pulmonary hemorrhage and Septic shock</b>	2	2.0
<b>Pulmonary hemorrhage and necrotizing enterocolitis</b>	1	1.0
<b>septic shock only</b>	9	9.0
<b>septic shock and acute kidney injury</b>	3	3.0
<b>Septic shock, necrotizing enterocolitis and acute kidney injury</b>	1	1.0
<b>Died</b>	49	49.0
<b>Discharged</b>	51	51.0

This table showed that the most common complication was septic shock either alone or with another complication then CNS infection (7%) followed by pulmonary hemorrhage (6%). Mortality was 51% of septic neonates

**Table (5): Results of blood culture in patient group**

<b>Culture</b>	<b>Frequency</b>	<b>Percent</b>
<b>No growth</b>	5	5%
<b>Gram negative</b>	70	70%
<b>Gram positive</b>	7	7%
<b>Mixed infection</b>	18	18%

The blood culture isolates showed that 95% of patient group had positive blood culture results with predominance of Gram negative organisms(70%)

**Table (6) Microbiological profile found in positive blood cultures of studied group**

<b>Blood culture</b>		
<b>Name of organism</b>	<b>Frequency</b>	<b>Percent</b>
<b>Klebsiella</b>	46	46.0
<b>Acinetobacter</b>	12	12.0
<b>Enterobacter</b>	6	6.0

<b>Pseudomonas</b>	5	5.0
<b>Atypical Gram -ve bacilli</b>	1	1.0
<b>Coagulase negative staphylococci</b>	4	4.0
<b>Entrococci</b>	2	2.0
<b>Methicillin resistant staphylococcus</b>	1	1.0
<b>Mixed infection</b>	18	18.0

<sup>2</sup> This table showed that the most frequent organism was *Klebsiella* (46%) followed by *Acinetobacter* (12%) and then *Enterobacter* (6%).

**Table(7) Result of tracheal aspirate in patient group**

<b>Tracheal aspirate culture</b>		
Name of organism	Frequency	Percent
<b>No growth</b>	20	20.0
<b>Klebsiella</b>	67	67.0
<b>Acinetobacter</b>	5	5.0
<b>Pseudomonas</b>	2	2.0
<b>Atypical Gram -ve bacilli</b>	1	1.0
<b>Gram -ve bacilli</b>	1	1.0
<b>Mixed infection</b>	2	2.0
<b>Escherichia coli</b>	1	1.0
<b>Candida albicans</b>	1	1.0

<sup>2</sup> This table showed that the most frequent organism was *Klebsiella* (67%) followed by *Acinetobacter* (5%)

**Table (8) Association between mortality and different characteristics of cases:**

characteristics	Alive (no=51)	Died (no=49)	P-value
Gestational age (mean±SD)	36.4±1.8	34.4±2.6	<0.001*
Birth weight	2.8±0.5	2.3±0.7	0.001*

<b>Gender</b>			
<b>Female</b>	13(35.1%)	24(64.9%)	0.015*
<b>Male</b>	38(60.3%)	25(39.7%)	
<b>Respiratory support</b>			
<b>Nasal oxygen</b>	16(100.0%)	0(0.0%)	
<b>Continues positive airway pressure</b>	26(92.9%)	2(7.1%)	
<b>Mechanical ventilation</b>	7(13.0%)	47(87.0%)	<0.001*
<b>Blood or blood elements transfusion</b>			
<b>Blood</b>	34(41.0%)	49(59.0%)	
<b>Plasma</b>	17(26.2%)	48(73.8%)	
<b>PLT</b>	5(12.8%)	34(87.2%)	<0.001*
<b>Need to inotropes</b>			
<b>No</b>	40(93.0%)	3(7.0%)	
<b>Yes</b>	11(19.3%)	46(80.7%)	<0.001*
<b>Respiratory Degree</b>			
<b>Complications of neonatal sepsis</b>			
<b>No</b>	47(100.0%)	0(0.0%)	
<b>Yes</b>	4(7.5%)	49(92.5%)	<0.001*
<b>Heart rate</b>	131.6±20.4	133.9±27.2	0.626
<b>Respiratory rate</b>	55.9±21.5	57.5±13.3	0.664
<b>systolic blood pressure</b>	75.3±6.9	71.3±9.5	0.020*
<b>Diastolic blood pressure</b>	45.8±6.7	41.9±7.7	0.009*

\* P-value is significant

This table showed that there was a significant association between mortality and younger gestational age, birth weight, female gender, need to mechanical ventilation, need to platelet transfusion, need to positive inotropic drugs, Complications in NICU, birth weight, and lower blood pressure.

## DISCUSSION

In the first month of life, when normally sterile fluids like blood or cerebrospinal fluid (CSF) harbor pathogenic microbes like bacteria, viruses, or fungi, a clinical syndrome known as neonatal sepsis can develop, causing hemodynamic abnormalities and other systemic symptoms [7]. It significantly impacts neurodevelopmental problems and infant mortality rates [8]. Symptoms, causes, and prognoses of infant sepsis were the foci of this investigation. Our study evaluated a total of 100 infants, 24 of whom were born naturally vaginally and 76 through cesarean surgery. Consistent with previous research, we have found the following: Researchers Seliem and Sultan [9] and Bates et al. [10] found that sepsis was more common in babies born by cesarean section than in vaginal births. On the other hand, compared to babies born by cesarean section, those born vaginally were more likely to develop sepsis [11]. After cesarean sections, neonatal sepsis rates significantly decreased, potentially due to the effectiveness of sterilization and intrapartum chemoprophylaxis.

Hamam et al. [12] found that neonates with neonatal sepsis had an average birth weight of  $2.309 \pm 0.770$  kg, which is in agreement with our patients' average weight of  $2.6 \pm 0.7$  kg. While Smertka M et al. [13] did not find a connection between low birth weight and an increased risk of sepsis, Schrag [14] did. Seventy percent of the women in our study had no history of problems during pregnancy, fourteen percent experienced premature membrane rupture (defined as occurring within the first eighteen hours), and two percent showed signs of chorioamnionitis. Only 8 cases showed pre-eclampsia and pregnancy-induced hypertension. Maternal illness in the form of gestational diabetes mellitus affected four cases (four percent), while obstructed labor affected five percent of the deliveries. That is in line with the findings of the 2014 study by Idris AA et al., [15] which found that 22.2% of the mothers whose babies had sepsis also had PROM. Conversely, Abdou et al. [16] found that 51.6% of the assessed pregnant women had no history of pregnancy complications. Of those, 8.9% had obstructed labor, 7.8% had eclampsia or pregnancy-induced hypertension, and 3.8% had PROM. The same percentage of women also had chorioamnionitis.

Out of 100 case, 55 case referred from another hospital, 35 were born at Beni-Suef university Hospital, 10 were admitted from emergency department at Beni-Suef university Hospital, against our results Ingale et al., [17] (found that (74.5%) of the infected babies were inborn. Regarding onset of neonatal sepsis 76% of our patients had

late onset sepsis, 24% had early onset sepsis, the same results found by Gaballah et al., 2022 [18] 457 out of 2,400 newborns (19%) positively recognized the prevalence of sepsis. Gosalia et al. [19] found that 181 patients (39.6%) had early onset newborn sepsis (EOS), while 276 cases (60.4%) had late onset neonatal sepsis (LOS). One possible explanation for the high number of LOS cases is the high number of invasive operations that preterm and low birth weight babies undergo, including endotracheal intubation and intravenous catheter placement. Abdelaziz et al. [20] found the opposite: early-onset sepsis in 130 patients (64.4%) and late-onset sepsis in 72 instances (35.6%). Based on the findings of this study, respiratory distress accounted for 90% of NICU admissions, either alone or in combination with other reasons. Convulsions accounted for 5% and jaundice for 4%.

The most common symptoms of neonatal sepsis, according to Shehab et al. [21], are difficulty breathing (54.6%), yellowing of the skin and eyes (4.9%), fluctuating body temperature (2.3%), and seizures (1.6%). Consistent with previous research [22], the most common cause for NICU hospitalizations (63.4% of cases) was respiratory distress, which includes hyaline membrane disorders, meconium aspiration syndrome, and fetal asphyxia. Shortly after, congenital cardiac disease was found in 9.5% of cases, and hypoglycemia was present in 4.8%. Also in agreement with these findings is Abdou et al. [16]. We found that respiratory distress (at 88% of patients), poor eating (73% of patients), and jaundice were the most common clinical symptoms of newborn sepsis. This confirms what West and Tabansi [23] found; they identified fever (26.6% of cases), poor sucking (22.5% of cases), jaundice (14.2%), and respiratory distress (30.2% of cases) as the most prevalent clinical signs of septicemia. Research by Basu and Bandyopadhyay [24] and Lakhey and Shakya [25] confirms that respiratory distress is the most common symptom of sepsis.

Our findings contradicted previous findings. The main symptoms of neonatal sepsis, according to Verma et al. [26] and Shalaby et al. [27], include a lack of interest in eating and insufficient suctioning.

According to our data, septic shock is the most common occurrence, either alone or in conjunction with another problem; second most common were central nervous system infections (7% of cases) and pulmonary bleeding (6% of cases). On the other hand, Al-Matary et al. [28], conducted a study in 2019. In 126 instances (59.4%), the most frequent outcome of neonatal sepsis was chronic lung disease (CLD). In 113 cases (53.3%), the second most common consequence was a longer hospital stay that exceeded 120 days. In 63 cases (23.8%), the third most common consequence was necrotizing enterocolitis (NEC).

Supporting the results of Jumah DS [29], this study found that while 49% of the infants tested died, 51% showed improvement and were discharged from the NICU. There was a 44.2% significant mortality rate among the 29 individuals. Also, in 2019, Hamam et al. [12], documented mortality rates hover around 57.24%.

On the other hand, a study that looked at the <sup>12</sup> pattern of microbial illnesses in the first 72 hours of a newborn's life in the NICU at Al Khafji General Hospital in Saudi Arabia found that only 12.2% of those babies died from infections [30]. Additionally, Ingale et al., [17]. reported a lower death rate (20.7%) among infected infants. Blood cultures most commonly contained Klebsiella (46%), Acinetobacter (12%), and Enterobacter (6%). In a similar vein, Gaballah et al. [18] carried out studies in Egypt. Klebsiella emerged as the most common pathogen, accounting for 54 percent (18 isolates). Hamam et al. [12]. reported that Klebsiella caused 22.89% of the 145 instances of sepsis at Tanta University Hospital in Egypt. According to Utomo et al. [31], Klebsiella is the primary pathogen in neonatal sepsis, as confirmed by culture [31], in Indonesia and other countries that use it [32,33]. Eltaib L and <sup>35</sup> Alshammari HA [30] discovered that Pseudomonas aeruginosa (13.5%) and other gram-negative bacteria were the main pathogens responsible for illnesses in infants, which contradicts our results. Additionally, in 2019, Thapa <sup>g</sup> and Sapkota [34] conducted research in Nepal. Acinetobacter species (32.1%) and Staphylococcus aureus (19.6%) were the most frequently isolated bacteria.

## CONCLUSION

<sup>28</sup> Among neonates who admitted at Beni-Suef university hospital with neonatal sepsis Sepsis was <sup>1</sup> more common in babies born by cesarean section than in <sup>32</sup> vaginal births .The majority of septic neonates were referred to our unit from other hospital, the most common cause of admission was respiratory distress, while the most common isolated organism was klebsiella. Septic shock was the main complication leading to death. The survival incidence was 51%. Prematurity, low birth weight, female gender, mechanical ventilation, platelet transfusion, Complications in neonatal intensive care unit during staying , intravenous infusion of positive inotropic drugs, and lower blood pressure were considered risk factors for death .AS there were significant associations between these parameters and mortality

## REFERENCES

1.Kimberlin DW. Red Book: 2018-2021 report of the committee on infectious diseases. 2018.doi:10.1542/9781610021470

- 2.** Wang D, Yin Y, Yao Y. Advances in sepsis-associated liver dysfunction. *Burns & trauma*. 2014 Jul;2(3):2321-3868.doi:10.4103/2321-3868.132689
- 3.** Klinger G, Levy I, Sirota L, Boyko V, Reichman B, Lerner-Geva L, Israel Neonatal Network. Epidemiology and risk factors for early onset sepsis among very-low-birthweight infants.*American journal of obstetrics and gynecology*. 2009 Jul 1; 201(1):38-e1. doi:10.1016/j.ajog.2009.03.006
- 4.** Yadav AK, Wilson CG, Prasad PL, Menon PK. Polymerase chain reaction in rapid diagnosis of neonatal sepsis. *Indian pediatrics*. 2005 Jul 1;42(7):681.doi:10.4103/0971-5916.178613
- 5.** Long SS, Pickering LK, Prober CG. Bacterial infections in the neonate. *Principles and Practice of Pediatric Infectious Diseases* (4th ed.). Elsevier. 2012.doi:10.1016/b978-1-4377-2702-9.00094-5
- 6.** Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA, Ehrenkranz RA, Lemons JA, Donovan EF, Stark AR, Tyson JE, Oh W. Late-onset sepsis in very low birth weight neonates: the experience of the NICHD Neonatal Research Network. *Pediatrics*. 2002 Aug 1;110(2):285-91.doi:10.1542/peds.110.2.285
- 7.** Shane AL, Sánchez PJ, Stoll BJ. Neonatal sepsis.*The lancet*. 2017 Oct 14;390(10104):1770-80. doi:10.1016/s0140-6736(17)31002-4
- 8.** Liang L, Kotadia N, English L, Kissoon N, Ansermino JM, Kabakyenga J, Lavoie PM, Wiens MO. Predictors of mortality in neonates and infants hospitalized with sepsis or serious infections in developing countries: a systematic review. *Frontiers in pediatrics*. 2018 Oct 4;6:277. doi:10.3389/fped.2018.00277

**9.** Seliem WA, Sultan AM. Etiology of early onset neonatal sepsis in neonatal intensive care unit–Mansoura, Egypt. *Journal of neonatal-perinatal medicine*. 2018 Jan 1; 11(3):323-30. doi:10.3233/npm-17128

**10** Bates M, Kabwe M, Zumla A. Neonatal sepsis and antibiotic resistance in developing countries. *The Pediatric Infectious Disease Journal*. 2014 Oct 1; 33(10):1097. doi:10.1097/inf.0000000000000388

**11.** Kardana IM. Incidence and factors associated with mortality of neonatal sepsis. *Paediatrica Indonesiana*. 2011 Jun 30; 51(3):144-8. doi:10.14238/pi51.3.2011.144-8

**12.** HAMAM SM, Shimaa M, Hamed M, Mohamed S. Incidence of neonatal sepsis and the causative organisms in neonatal intensive care unit of tanta university hospital. *The Medical Journal of Cairo University*. 2019 Dec 1; 87(December):5323-32. doi:10.21608/mjcu.2019.89803

**13.** Smertka M et al. Serum and urinary NGAL in septic newborns. *BioMed research international*. 2014; 2014(1):717318. doi:10.1155/2014/717318

**14.** Schrag SJ, Cutland CL, Zell ER, Kuwanda L, Buchmann EJ, Velaphi SC, Groome MJ, Madhi SA, PoPS Trial Team. Risk factors for neonatal sepsis and perinatal death among infants enrolled in the prevention of perinatal sepsis trial, Soweto, South Africa. *The Pediatric infectious disease journal*. 2012 Aug 1; 31(8):821-6. doi:10.1097/inf.0b013e31825c4b5a



**15.**Idris AA, Tosson AM, Hassan RM, Abdelkader SL. Neonatal sepsis and antibiotic susceptibility in NICUs of Cairo University hospitals *Journal of Arab Child*. 2014 Dec;319(2126):1-1.doi:10.12816/0014277

**16.**Abdou AA, El-Latef A, Amal M, Elnaggar MM. Outcome of Neonatal Sepsis in Neonatal Intensive Care Unit in Zagazig University Hospitals. *The Egyptian Journal of Hospital Medicine*. 2022 Apr 1;87(1):1899-906. doi:10.21608/ejhm.2022.231653

**17.**Ingale HD, Kongre VA, Bharadwaj RS. A study of infections in neonatal intensive care unit at a tertiary care hospital. *Int J ContempPediatr*. 2017 Jul; 4(4):1349-56.doi:10.18203/2349-3291.ijcp20172664

**18.**Gaballah AH, Shawky S, Amer AN. Microbiological profiles of neonatal sepsis in northern Egypt. *Microbes and Infectious Diseases*. 2022 Aug 1;3(3):645-56. doi:10.21608/mid.2022.129600.1265

**19.**Gosalia E, Mistry M, Goswami Y, Gosalia V, Vasa P. A bacteriological profile of neonatal septicemia (Study in Tertiary Care Hospital, Rajkot). *NJIRM*. 2013;4(2):44-7. doi:10.70284/njirm.v4i2.2146

**20.**Abdelaziz M, Hamad Alnil Y, Hashim O, Bashir T, Mahjoub ES. Microbiological profile of neonatal sepsis at a maternity hospital in Omdurman, Sudan. *Sudan Journal of Medical Sciences*. 2019 Jul 22; 14(1):45-51. doi:10.18502/156282

**21.**Shehab El-Din EM, El-Sokkary MM, Bassiouny MR, Hassan R. Epidemiology of neonatal sepsis and implicated pathogens: a study from Egypt. *BioMed research international*. 2015;2015(1):509484. doi:10.1155/2015/509484

**22.** Mehar V, Yadav D, Somani P, Bhatambare G, Mulye S, Singh K. Neonatal sepsis in a tertiary care center in central India: microbiological profile, antimicrobial sensitivity pattern and outcome. *Journal of Neonatal-Perinatal Medicine*. 2013 Jan 1; 6(2):165-72. doi:10.3233/npm-1367312

**23.** West BA, Tabansi PN. Prevalence of neonatal septicaemia in the University of port harcourt teaching hospital, Nigeria. *Nigerian Journal of Paediatrics*. 2014; 41(1):33-7.  
doi:10.4314/njp.v41i1.6

**24.** Basu R, Bandyopadhyay S. Study on correlation between sepsis screening and blood culture in neonatal sepsis. *J Dent Med Sci*. 2014 May; 13(5):52â. doi:10.9790/0853-13555256

**25.** Lakhey A, Shakya H. Role of sepsis screening in early diagnosis of neonatal sepsis. *Journal of Pathology of Nepal*. 2017 Mar 30;7(1):1103-10.  
doi:10.3126/jpn.v7i1.16944

**26.** Verma P, Berwal PK, Nagaraj N, Swami S, Jivaji P, Narayan S. Neonatal sepsis: epidemiology, clinical spectrum, recent antimicrobial agents and their antibiotic susceptibility pattern. *Int J ContempPediater*. 2015 Jul;2(3):176-80. doi:10.18203/2349-3291.ijcp20150523

**27.** Shalaby MM, Sobeih AA, Abdulghany WE, Behiry EG, Ismail YM, Abd-El-Aziz MA. Mean platelet volume and serum uric acid in neonatal

sepsis: A case-control study. *Annals of medicine and surgery*. 2017 Aug 1;20:97-102. doi:10.1016/j.amsu.2017.06.015

**28.** Al-Matary A, Heena H, AlSarheed AS, Ouda W, AlShahrani DA, Wani TA, Qaraqei M, Abu-Shaheen A. Characteristics of neonatal Sepsis at a tertiary care hospital in Saudi Arabia. *Journal of infection and public health*. 2019 Sep 1;12(5):666-72. doi:10.1016/j.jiph.2019.03.007

**29.** Jumah DS. Predictors of mortality outcome in neonatal sepsis. *The Medical Journal of Basrah University*. 2007; 25(1). doi:10.33762/mjbu.2007.48118

**30.** Eltaib L, Alshammari HA. Pattern of Microbial Infections during the First 72 Hours of Neonate Life at Khafji General Hospital Neonatal Intensive Care Unit. *Asian Journal of Pharmaceutical Research and Health Care*. 2020:189-97. doi:10.18311/ajprhc/2020/25739

**31.** Utomo MT, Sumitro KR, Etika R, Widodo AD. Current-proven neonatal sepsis in Indonesian tertiary neonatal intensive care unit: a hematological and microbiological profile. *Iranian Journal of Microbiology*. 2021 Jun;13(3):266. doi:10.18502/ijm.v13i3.6386

**32.** Gago C, Alexandre S, Marçal M & Tuna M. Neonatal Septic Shock in a Neonatal Intensive Care Unit-Trends In Incidence and Therapeutic Challenges. *J Neonatol Clin Pediatr*. 2022; 9: 091. Of: 5:2.

doi:10.24966/ncp-878x/100091

**33.** Dedeke I, Arowosegbe A, Shittu O, Ojo D, Akingbade O. Neonatal sepsis in a Nigerian tertiary hospital: clinical features, clinical outcome, aetiology and antibiotic susceptibility pattern. *Southern African Journal of Infectious Diseases*. 2017 Dec 1;32(4):127-31. doi:10.1080/23120053.2017.1335962

**34.**Thapa S, Sapkota LB. Changing trend of neonatal septicemia and antibiotic susceptibility pattern of isolates in Nepal. *International journal of pediatrics*. 2019;2019(1):3784529. doi:10.1155/2019/3784529