

The significance of follow-up in patients with dysmorphic features: a case from clinical practice

By Mariya Levkova

TYPE OF ARTICLE: Case Report

The significance of follow-up in patients with dysmorphic features: a case from clinical practice

Mariya Levkova^{1,2}, Milena Stoyanova^{1,2}, Mari Hachmeriyan^{1,2}

³
¹ Department of Medical Genetics, Medical University Varna, Marin Drinov Str 55, Varna, Bulgaria

² Laboratory of Medical Genetics, St. Marina Hospital, Hristo Smirnenski Blv 1, Varna, Bulgaria

corresponding author

¹²
e-mail: maria.levkova171@gmail.com

ABSTRACT

⁴
Background. Kabuki syndrome is a rare disorder, that is characterized by typical facial dysmorphism, hypotonia, delay in intellectual and motor development.

⁵
Case Report. We present a case of a girl in whom polycystic left kidney was prenatally established. Born prematurely in 37 weeks by C section due to oligohydramnios. After birth, atresia of the anus with fistula, cysts in the left and reduced dimensions of the right kidney, were further established. A normal female karyotype was found and targeted sequencing analysis was conducted on a panel of 81 genes associated with renal abnormalities – no pathogenic variants were detected. The child was then followed up by its general practitioner. ⁴ At the age of 2 years, she was again referred for genetic counseling, which revealed ¹³ the following dysmorphic signs – long eye palpebral fissuras with ectropion of the lower eyelid, sparsed lateral eyebrows, depression of the nasal bridge,

brachydactyly and others characteristic of Kabuki's syndrome. The conducted molecular genetic analysis confirmed the clinical diagnosis – a likely pathogenic variant in the KMT2D gene was established.

Conclusions. Certain pathognomonic facial may not present at birth and only appear after a few years. Therefore, monitoring of the evolution of dysmorphic traits is required.

Keywords: Kabuki syndrome, KMT2D, dysmorphic features, genetic counseling

Abbreviations:

KS – Kabuki syndrome

Gr - grams

Cm - centimetres

IQ - intellectual quotient

INTRODUCTION

The detection of a certain genetic disorder depends both on the physical examination and the molecular-genetic analysis. Approximately 30–40% of these conditions are associated with specific dysmorphic features, and some of them may be highly indicative of a specific disorder [1].

However, the typical dysmorphic traits may not be present at birth and develop later in life, which makes it difficult to diagnose a genetic disorder right away from the first patient's visit.

Therefore, it is not surprising that the average time for diagnosing a patient with a rare disorder is around 6 years from the first symptoms, and in the meantime, the patient would receive several wrong diagnoses [2].

Nevertheless, an early diagnosis is crucial since it makes it possible to start screening for the underlying condition's related issues, such as cardiovascular or malignant ones [3].

¹¹ Kabuki syndrome (KS) is a rare disorder, which was first reported in Japan in 1980s [4]. It was named like this because of the distinctive facial features that mimic the makeup of performers in Japanese Kabuki theater [4]. ⁹ It is caused by pathogenic variants in the genes *KMT2D* or *KDM6A* and its prevalence is around 1 in 32 000 [5]. The typical dysmorphic features of KS are ¹ long palpebral fissures with eversion of the lateral third of the lower eyelid and two or more of the following: arched and broad eyebrows with the lateral third displaying notching or sparseness; short columella with depressed nasal tip; large, prominent, or cupped ears, and persistent fingertip pads [6, 7]. ² Additional symptoms are short stature, microcephaly, cleft palate, lip pits, hearing loss, congenital heart defects, feeding difficulties and immunological disorders [4,7].

Despite the striking facial features, typical for KS, the diagnosis might be delayed because these features develop in the first several years of life [4].

⁵ We present a case of a patient with KS and the course of the diagnostic process.

CASE REPORT

⁸ The patient is a 2-year-old girl born from first pregnancy. A polycystic kidney and one cyst in the other kidney were noted in the 35th gestational week. The girl was born prematurely per C section in the 37th week due to oligohydramnios, weight – 3200 gr, length – 50 cm. Anal atresia with a fistula, multiple cysts in the left kidney, decreased size of the right kidney were noted after the birth of the baby. The girl was clinically diagnosed with autosomal recessive polycystic kidney disease and was referred to genetic counseling. The karyotype of the girl was normal – 46, XX. Targeted sequencing for 81 genes, associated with kidney anomalies and polycystic kidney disease was performed, but ⁷ no pathogenic or likely pathogenic variants were reported. After that, the child was followed by its general practitioner.

⁷ At two years of age, the girl was again referred to genetic counseling due to the presence of dysmorphic features and intellectual disability. The girl presented with ¹ long palpebral fissures with eversion of the lower eyelid, sparse lateral eyebrows, depressed nasal bridge, anteverted nares, ⁸ low set ears with overfolded helix and linear creases of the left earlobe, high-arched palate, brachydactyly, ulnar deviation of both thumbs, and persistent fingertip pads (Figure 1). The

suspected diagnosis was Kabuki syndrome, based on the similarity with the cardinal features of KS.

Targeted sequencing of 1902 genes reported a likely pathogenic variant p.(Arg1709Hisfs*25) in *KMT2D* gene, which was associated with KS. In our case there was no family history of other affected members and the pathogenic variant occurred de novo.

The child was referred to a pediatric department, specialized in patients with rare disorders and is regularly screened for the manifestation of additional symptoms, typical for KS.

DISCUSSION

KS is a heterogenous disorder, which involves various systems. The phenotypic features vary over time and the cardinal dysmorphism may manifest later in life, like it was in our case. That is why the patients should be reevaluated for the presence of specific dysmorphic features in certain time intervals with the application of the international ¹ diagnostic criteria for KS [7]. However, molecular-genetic testing is required to further prove the suspected underlying condition.

The intellectual disability is typical for this condition and the intellectual quotient (IQ) of the patients is usually below 70, but there are also cases described of mild and moderate intellectual disability [8]. Infantile hypotonia and epilepsy are also reported in ¹⁴ patients with Kabuki syndrome [7,8].

A ¹⁰ congenital heart defect affects about 80% of people with KS, the most common being atrial septal defect, ventricular septal defect and coarctation of aorta [4]. Patients with KS typically present with poor feeding, which may be due to severe reflux. Chronic diarrhea is also described [4]. As was the case in our instance, anorectal anomalies such as atresia of the anus are possible. Involvement of the kidneys is typical for KS, however, cysts are not a typical finding (described in our case). More than 40% of afflicted people with KS have present with hydronephrosis, renal dysplasia, and horseshoe kidney [4].

Endocrine issues include hypothyroidism, short stature with unknown etiology, obesity, cryptorchidism in boys are described [8, 9]. KS is associated with immune dysfunction. This could be explained by the fact that the responsible genes *KMT2D* and *KDM6A* participate in B-lymphocyte differentiation. Hypogammaglobulinemia is a typical finding and patients with KS

may present later with common variable immune deficiency. There is also an increased risk of autoimmune disorders such as hemolytic anemia, vitiligo, and immune thrombocytopenia [4,8]. KS may also predispose to malignant diseases and the patients should be regularly screened for such complications [4].

Over one-third of the affected individuals have ocular abnormalities, such as blue sclerae, strabismus, nystagmus, nocturnal lagophthalmos and dry eye, refractive error, and coloboma [4]. Hearing loss affects up to 50% of affected individuals, and it may be linked to chronic otitis media [4]. Numerous dental anomalies are also described in more than 60% of KS patients, most frequently hypodontia, widely spaced teeth, irregularly formed teeth, missing lateral, upper, and lower incisors, and malocclusion [4].

CONCLUSION

Some of the dysmorphic syndromes are characterized by typical facial features, which are pathognomonic. However, it is possible that they are not present at birth, but develop after a few years. This requires sustainable monitoring of undiagnosed patients and, if necessary, reanalysis of the clinical diagnosis. Thorough follow-up of phenotypic traits and their documentation are key in the evaluation of patients with rare dysmorphic traits.

PATIENT CONSENT

Informed consent was obtained from legal guardians.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

AUTHOR'S CONTRIBUTIONS

Conceptualization, M.L. and M.S.; investigation, M.L. and M.S.; resources, M.H.; writing—original draft preparation, M.L.; writing—review and editing, M.S. and M.H. All authors have read and agreed to the published version of the manuscript.

ACKNOWLEDGEMENTS

Not applicable.

REFERENCES

1. Reardon W, Donnai D. Dysmorphology demystified. *Archives of Disease in Childhood-Fetal and Neonatal Edition*. 2007;92(3):F225-F9.
2. Blöß S, Klemann C, Rother A-K, Mehmecke S, Schumacher U, Mücke U, et al. Diagnostic needs for rare diseases and shared prediagnostic phenomena: Results of a German-wide expert Delphi survey. *PLoS One*. 2017;12(2):e0172532.
3. Geremek M, Szklanny K. Deep learning-based analysis of face images as a screening tool for genetic syndromes. *Sensors*. 2021;21(19):6595.
4. Boniel S, Szymańska K, Śmigiel R, Szczałuba K. Kabuki Syndrome—Clinical Review with Molecular Aspects. *Genes*, 12 (4), 468. 2021.
5. Cheon C-K, Ko JM. Kabuki syndrome: clinical and molecular characteristics. *Korean journal of pediatrics*. 2015;58(9):317.
6. Adam MP, Hudgins L, Hannibal M. Kabuki syndrome. 2021.
7. Adam MP, Banka S, Bjornsson HT, Bodamer O, Chudley AE, Harris J, et al. Kabuki syndrome: international consensus diagnostic criteria. *Journal of medical genetics*. 2019;56(2):89-95.
8. Di Candia F, Fontana P, Paglia P, Falco M, Rosano C, Piscopo C, et al. Clinical heterogeneity of Kabuki syndrome in a cohort of Italian patients and review of the literature. *European journal of pediatrics*. 2022;181(1):171-87.
9. van Montfort L, Gerver WJM, Kooger BL, Plat J, Bierau J, Stumpel CT, et al. Follow-up study of growth hormone therapy in children with Kabuki syndrome: two-year treatment results. *Hormone Research in Paediatrics*. 2021;94(7-8):285-96.

FIGURES



Figure 1. Evolution of the dysmorphic traits over time A) Newborn B) Two-months old C) Two-year old